

ON MULTIPLICITY AND FREE ABSORPTION FOR FREE ARAKI-WOODS FACTORS.

DIMITRI SHLYAKHTENKO

ABSTRACT. We show that Ozawa's recent results on solid von Neumann algebras imply that there are free Araki-Woods factors, which fail to have free absorption. We also show that a free Araki-Woods factors $\Gamma(\mu, n)$ associated to a measure and a multiplicity function n may non-trivially depend on the multiplicity function.

In [6] we have associated to the absolute continuity class of a symmetric measure μ on $\mathbb{R}$, $\mu(X) = \mu(-X)$, and a multiplicity function $n : \mathbb{R} \rightarrow \mathbb{N} \cup \{+\infty\}$, $n(x) = n(-x)$, a von Neumann algebra denoted by $\Gamma(\mu, n)$ ¹. This construction is the free probability analog of the Araki-Woods construction of hyperfinite type III factors using quasi-free states [1, 2]. The algebras $\Gamma(\mu, n)$ are called free Araki-Woods factors; they are factors of type III if μ is not concentrated on the set $\{0\}$. The complete classification of free Araki-Woods factors remains an interesting outstanding problem, and was achieved in [6] only for atomic measures μ . In our recent work [5] we have introduced new invariants for type III factors, which show that the algebras $\Gamma(\mu, n)$ depend to a large extent on the absolute continuity class of the measure μ . On the other hand, it remained completely unknown in general whether the multiplicity function n matters at all. In the case that μ is atomic and not concentrated on $\{0\}$, we have shown in [6] that $\Gamma(\mu, n)$ is independent of the multiplicity function, leading one to wonder if the same holds in general.

Our construction of a free Araki-Woods factor $\Gamma(\mu, n)$ produces also a state $\phi_{\mu, n} : \Gamma(\mu, n) \rightarrow \mathbb{C}$, which is the analog of the quasi-free state in the classical Araki-Woods construction. We have shown in [6] that the reduced free product of free Araki-Woods factors is given by:

$$(\Gamma(\mu, n), \phi_{\mu, n}) * (\Gamma(\mu', n'), \phi_{\mu', n'}) \cong (\Gamma(\mu + \mu', n + n'), \phi_{\mu + \mu', n + n'}),$$

with the isomorphism preserving the indicated states.

Independence of n then says in particular that

$$(\Gamma(\mu, n), \phi_{\mu, n}) * (\Gamma(\mu, n), \phi_{\mu, n}) \cong \Gamma(\mu, n)$$

(via an isomorphism that does not necessarily take the free product state to $\phi_{\mu, n}$).

Date: 8th February 2008.

This material is based upon work performed for the Clay Mathematics Institute.

¹Our original notation for this algebra is $\Gamma(\mathcal{H}_{\mathbb{R}}, U_t)''$, where $t \mapsto U_t$ is an orthogonal representation of $\mathbb{R}$ on a real Hilbert space. This data is of course completely equivalent to specifying the spectral measure μ and multiplicity function n of such a representation. It is more convenient to use the notation $\Gamma(\mu, n)$ in the present paper.

Another property of free Araki-Woods factors which holds for μ atomic and not concentrated on $\{0\}$ is the “free absorption” property:

$$\Gamma(\mu, n) \cong (\Gamma(\mu, n), \phi_{\mu, n}) * (L(\mathbb{F}_\infty), \tau),$$

where $L(\mathbb{F}_\infty)$ denotes the von Neumann algebra of a free group on an infinite number of generators.

In this note we rely on the amazing recent result of Ozawa [3] showing that von Neumann algebras of hyperbolic groups (in particular, of free groups) are “solid”: the relative commutant of any diffuse unital subalgebra is hyperfinite. From this result, we derive the following theorem, which shows that in the case that μ is non-atomic, free absorption and independence of the multiplicity function need not hold.

Theorem 1. *Let λ denote the Lebesgue measure on $\mathbb{R}$, and let δ_0 denote the delta measure at 0. Then:*

- (a) $\Gamma(\lambda + \delta_0, 1) \not\cong \Gamma(\lambda + \delta_0, 2)$. Thus $\Gamma(\lambda + \delta_0, n)$ depends on n .
- (b) $\Gamma(\lambda, 1) \not\cong (\Gamma(\lambda, 1), \phi_{\lambda, 1}) * (L(\mathbb{F}_\infty), \tau)$. Thus free absorption fails.

Proof. It is known that the core of $\Gamma(\lambda, 1)$ is isomorphic to $L(\mathbb{F}_\infty) \otimes B(\ell^2)$ (see [7]). We’ll first show that the core of $\Gamma(\lambda + \delta_0, 1)$ is isomorphic to $L(\mathbb{F}_\infty) \otimes B(\ell^2)$.

We have that $(\Gamma(\lambda + \delta_0, 1), \phi_{\lambda + \delta_0, 1}) \cong (\Gamma(\lambda, 1), \phi_{\lambda, 1}) * (\Gamma(\delta_0, 1), \phi_{\delta_0, 1})$. Thus by [7], the core of $\Gamma(\lambda + \delta_0, 1)$ is isomorphic to the amalgamated free product

$$(\Gamma(\lambda, 1) \rtimes_{\sigma^{\phi_{\lambda, 1}}} \mathbb{R}, E_1) *_{L(\mathbb{R})} (\Gamma(\delta_0, 1) \rtimes_{\sigma^{\phi_{\delta_0, 1}}} \mathbb{R}, E_2).$$

Here E_j are the conditional expectations onto the group von Neumann algebra of $\mathbb{R}$ (viewed as a subalgebra of the crossed product), associated to the states $\phi_{\lambda, 1}$ and $\phi_{\delta_0, 1}$, respectively.

Since $\Gamma(\delta_0, 1) \cong L^\infty[0, 1]$ by [6], we have that the core of $\Gamma(\lambda + \delta_0, 1)$ is isomorphic to

$$(\Gamma(\lambda, 1) \rtimes_{\sigma^{\phi_{\lambda, 1}}} \mathbb{R}, E_1) *_{L(\mathbb{R})} (L(\mathbb{R}) \otimes A, \text{id} \otimes \tau),$$

where $A \cong L^\infty[0, 1]$, and τ is some normal faithful trace on A .

Let $\text{Tr} : L(\mathbb{R}) \rightarrow \mathbb{C}$ be the semifinite trace associated to integration against Lebesgue measure in the identification $L(\mathbb{R}) \cong L^\infty(\mathbb{R})$. Then by [8, 4], we have that $(\Gamma(\lambda, 1) \rtimes \mathbb{R}, E_1)$ is isomorphic to $(\Phi(L(\mathbb{R}), \text{Tr}), E_{L(\mathbb{R})})$ (see [4] for this notation). In other words, the core of $\Gamma(\lambda, 1)$, when viewed as a non-commutative probability space over $L(\mathbb{R})$ is generated by $L(\mathbb{R})$ and an operator-valued semicircular variable X of covariance $\eta : L^\infty(\mathbb{R}) \rightarrow L^\infty(\mathbb{R})$, where η is a completely-positive map given by

$$\eta(f)(x) = \int f(y) \omega(x, y) dy,$$

where $\omega > 0$ is a certain function on $\mathbb{R}^2$ (i.e., $\omega(x, y) dx dy$ and $dx dy$ are mutually absolutely continuous).

Since the core of $\Gamma(\lambda + \delta_0, 1)$ is given as an amalgamated free product, it follows that X is free with amalgamation over $L(\mathbb{R})$ from $A \otimes L(\mathbb{R})$. Thus by [7], if we view X as an operator-valued random variable with values in $A \otimes L(\mathbb{R})$, it is still semicircular and has covariance $\eta \circ (\text{id} \otimes \tau) \sim \text{Tr} \circ (\text{id} \otimes \tau) : L(\mathbb{R}) \otimes A \rightarrow \mathbb{C}$. But $(L(\mathbb{R}) \otimes A, \text{Tr} \otimes (\text{id} \otimes \tau)) \cong (L(\mathbb{R}), \text{Tr})$. Thus the core of $\Gamma(\lambda + \delta_0, 1)$ is isomorphic to $\Phi(L(\mathbb{R}), \text{Tr}) \cong L(\mathbb{F}_\infty) \otimes B(\ell^2)$ by [7].

We now prove the non-isomorphisms claimed in (a) and (b). It is sufficient to show that the cores of $\Gamma(\lambda + \delta_0, 2)$ and $(\Gamma(\lambda, 1), \phi_{\lambda,1}) * (L(\mathbb{F}_\infty), \tau)$ are not isomorphic to $L(\mathbb{F}_\infty) \otimes B(\ell^2)$.

The core of $\Gamma(\lambda + \delta_0, 2)$ is isomorphic, as before, to

$$(\Gamma(\lambda, 2) \rtimes_{\sigma^{\phi_{\lambda,2}}} \mathbb{R}, E_3) *_{L(\mathbb{R})} (\Gamma(\delta_0, 2) \rtimes_{\sigma^{\phi_{\delta_0,2}}} \mathbb{R}, E_4).$$

But by [6], $(\Gamma(\delta_0, 2), \phi_{\delta_0,2}) \cong (L(\mathbb{F}_2), \tau)$, and the core of $\Gamma(\lambda + \delta_0, 2)$ is given by

$$(\Gamma(\lambda, 2) \rtimes_{\sigma^{\phi_{\lambda,2}}} \mathbb{R}, E_3) *_{L(\mathbb{R})} (L(\mathbb{F}_2) \otimes L(\mathbb{R}), \tau \otimes \text{id}).$$

Now choose an abelian diffuse subalgebra $B \subset L(\mathbb{R})$, so that the restriction of the trace on the core of $\Gamma(\lambda + \delta_0, 2)$ to B is finite. Denote by p the unit of B . Then the relative commutant of $B = pBp$ inside of $p(\Gamma(\lambda, 2) \rtimes_{\sigma^{\phi_{\lambda,2}}} \mathbb{R}, E_3) *_{L(\mathbb{R})} (L(\mathbb{F}_2) \otimes L(\mathbb{R}), \tau \otimes \text{id})p$ contains $L(\mathbb{F}_2)$ and so is not hyperfinite. Thus $p(\Gamma(\lambda, 2) \rtimes_{\sigma^{\phi_{\lambda,2}}} \mathbb{R}, E_3) *_{L(\mathbb{R})} (L(\mathbb{F}_2) \otimes L(\mathbb{R}), \tau \otimes \text{id})p$ is not solid. Thus the core of $\Gamma(\lambda + \delta_0, 2)$ is not a free group factor, by Ozawa's result [3]. This proves (a).

The core of $(\Gamma(\lambda, 1), \phi_{\lambda,1}) * (L(\mathbb{F}_\infty), \tau)$ is, as before, given by

$$(\Gamma(\lambda, 1) \rtimes_{\sigma^{\phi_{\lambda,1}}} \mathbb{R}, E_5) *_{L(\mathbb{R})} (L(\mathbb{F}_\infty) \otimes L(\mathbb{R}), \tau \otimes \text{id}).$$

Using exactly the same argument as before, we find that the core is not $L(\mathbb{F}_\infty) \otimes B(\ell^2)$. $\square$

Note that the core of $\Gamma(\lambda + \delta_0, 2)$ is related to the factor $L(\mathbb{F}_\infty) * (L(\mathbb{F}_2) \otimes L^\infty[0, 1])$ (the core is, in some loose sense, “ $(L(\mathbb{F}_\infty) \otimes B(\ell^2)) * (L(\mathbb{F}_2) \otimes L^\infty(\mathbb{R}))$ ”). This explains our interest in this factor (see the remark in [3]).

There is, however, a situation where free absorption holds, and the multiplicity function does not matter. For a measure ν and a number $\lambda \in \mathbb{R}$, write ν_λ for the translate of ν by λ , and ν^λ for $\nu_\lambda + \nu_{-\lambda}$.

Theorem 2. *Assume that $\mu = \mu_c + \mu_a$, where μ_a is atomic and μ_c has no atoms. Assume that $\mu_a \neq 0$, and let $\Gamma = \{\gamma_1, \gamma_2, \dots\}$ be the subgroup of $\mathbb{R}$ generated by its support. Assume that $\Gamma \neq \{0\}$. Let $\nu = \sum_j 2^{-j} \mu^{\gamma_j}$. Then there is a state-preserving isomorphism*

$$(\Gamma(\mu, n), \phi_{\mu,n}) \cong (\Gamma(\nu, \infty), \phi_{\nu,\infty}).$$

In particular, $\Gamma(\mu, n)$ does not depend on n , and free absorption holds for $(\Gamma(\mu, n), \phi_{\mu,n})$.

Proof. We first note that the matricial model argument in [6] implies that if ϕ is a normal state on $B(\ell^2)$ given by

$$\theta(T) = \text{Tr}(DT),$$

where D is a trace-class diagonal matrix with eigenvalues $\exp(-\lambda_1), \exp(-\lambda_2), \dots$, with $\lambda_1, \lambda_2, \dots$ non-negative, then for any ν and m ,

$$(B(\ell^2), \theta) * (\Gamma(\nu, m), \phi_{\nu,m}) \cong (\Gamma(\sum 2^{-j} \nu^{\lambda_j}, \infty) \otimes B(\ell^2), \phi_{\sum 2^{-j} \nu^{\lambda_j}, \infty} \otimes \theta).$$

Moreover, the isomorphism is state-preserving.

It follows that if the set $\{0\} \cup \{\pm \lambda_j : j = 1, 2, \dots\}$ is an additive group, then

$$\begin{aligned} (B(\ell^2), \theta) * (\Gamma(\sum 2^{-j} \nu^{\lambda_j}, \infty) &\cong (\Gamma(\sum 2^{-j} \nu^{\lambda_j}, \infty) \otimes B(\ell^2), \phi_{\sum 2^{-j} \nu^{\lambda_j}, \infty} \otimes \theta) \\ &\cong (B(\ell^2), \theta) * (\Gamma(\nu, m), \phi_{\nu,m}), \end{aligned}$$

via state-preserving isomorphisms.

Returning now to our situation, note that $(\Gamma(\mu, n), \phi_{\mu, n}) \cong (\Gamma(\mu_c, n), \phi_{\mu_c, n}) * (\Gamma(\mu_a, n), \phi_{\mu_a, n})$. By [6], $(\Gamma(\mu_a, n), \phi_{\mu_a, n}) \cong (\Gamma(\mu_a, n), \phi(\mu_a, n)) * (B(\ell^2), \theta)$, where $\theta(T) = \text{Tr}(DT)$, and D is a diagonal operator with eigenvalues $\{e^{\gamma_j} : \gamma_j < 0\}$. Thus $(\Gamma(\mu, n), \phi_{\mu, n}) \cong (\Gamma(\mu, n), \phi_{\mu, n}) * (B(\ell^2), \theta)$. By the argument above, it follows that

$$\begin{aligned} (\Gamma(\mu, n), \phi_{\mu, n}) &\cong (\Gamma(\mu, n), \phi_{\mu, n}) * (B(\ell^2), \theta) \\ &\cong (\Gamma(\sum 2^{-j} \mu^{\gamma_j}, \infty), \phi_{\sum 2^{-j} \mu^{\gamma_j}, \infty}) * (B(\ell^2), \theta) \\ &\cong (\Gamma(\sum 2^{-j} \mu^{\gamma_j}, \infty), \phi_{\sum 2^{-j} \mu^{\gamma_j}, \infty}). \end{aligned}$$

Since $\sum 2^{-j} \mu^{\gamma_j}$ has an atom at zero, it follows that

$$\begin{aligned} (\Gamma(\mu, n), \phi_{\mu, n}) &\cong (\Gamma(\sum 2^{-j} \mu^{\gamma_j}, \infty), \phi_{\sum 2^{-j} \mu^{\gamma_j}, \infty}) \\ &\cong (\Gamma(\sum 2^{-j} \mu^{\gamma_j}, \infty), \phi_{\sum 2^{-j} \mu^{\gamma_j}, \infty}) * (L(\mathbb{F}_\infty), \tau) \\ &\cong (\Gamma(\mu, n), \phi_{\mu, n}) * (L(\mathbb{F}_\infty), \tau), \end{aligned}$$

so that free absorption holds. □

REFERENCES

1. H. Araki, *On quasifree states of CAR and Bogoliubov automorphisms.*, Publ. RIMS Kyoto Univ. **6** (1970), 385–442.
2. H. Araki and E. J. Woods, *A classification of factors*, Publ. RIMS Kyoto Univ. series A **4** (1968), 51–130.
3. N. Ozawa, *Solid von Neumann algebras*, Preprint, 2003.
4. S. Popa and D. Shlyakhtenko, *Universal properties of $L(\mathbb{F}_\infty)$ in subfactor theory*, MSRI preprint 2000-032, to appear in Acta Math., 2000.
5. D. Shlyakhtenko, *On the classification of full factors of type III*, Preprint, 2002.
6. ———, *Free quasi-free states*, Pacific J. Math **177** (1997), 329–368.
7. ———, *Some applications of freeness with amalgamation*, J. reine angew. Math. **500** (1998), 191–212.
8. ———, *A-valued semicircular systems*, J. Func. Anal **166** (1999), 1–47.

DEPARTMENT OF MATHEMATICS, UCLA, LOS ANGELES, CA 90095
E-mail address: shlyakht@math.ucla.edu